

OVACE® PLUS FOAM


(sodium sulfacetamide 9.8%)

Rx Only

FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.

DESCRIPTION: Each gram contains 98 mg of sodium sulfacetamide in a vehicle consisting of: benzyl alcohol, cetyl alcohol, fragrance, glyceryl stearate (and) PEG-100 stearate, magnesium aluminum silicate, phenoxyethanol, propylene glycol, purified water, sodium lauryl sulfate, sodium magnesium silicate, sodium thiosulfate, stearyl alcohol, and xanthan gum. This product also contains tetrafluoroethane (propellant).

Sodium sulfacetamide is a sulfonamide with antibacterial activity while sulfur acts as a keratolytic agent. Sodium sulfacetamide is $C_8H_9N_2NaO_3 \cdot H_2O$ with molecular weight of 254.24. Chemically, sodium sulfacetamide is N-[(4-aminophenyl)sulfonyl]-acetamide, monosodium salt, monohydrate. The structural formula is:



Sodium sulfacetamide is an odorless, white, crystalline powder with a bitter taste. It is freely soluble in water, sparingly soluble in alcohol, while practically insoluble in benzene, in chloroform and in ether.

CLINICAL PHARMACOLOGY: Sodium sulfacetamide exerts a bacteriostatic effect against sulfonamide sensitive Gram-positive and Gram-negative microorganisms commonly isolated from secondary cutaneous pyogenic infections. It acts by restricting the synthesis of folic acid required by bacteria for growth, by its competition with para-aminobenzoic acid. There is no clinical data available on the degree and rate of systemic absorption of this product when applied to the skin or scalp. However, significant absorption of sodium sulfacetamide through the skin has been reported.

The following in vitro data is available but the clinical significance is unknown. Organisms that show susceptibility to sodium sulfacetamide are: Streptococci, Staphylococci, E. coli, Klebsiella pneumoniae, Pseudomonas pyocyanea, Salmonella species, Proteus vulgaris, Nocardia and Actinomyces.

INDICATIONS: This product is intended for topical application in the following scaling dermatoses: seborrheic dermatitis and seborrhea sicca (dandruff). It also is indicated for the treatment of secondary bacterial infections of the skin due to organisms susceptible to sulfonamides.

CONTRAINDICATIONS: This product is contraindicated in persons with known or suspected hypersensitivity to any of the ingredients of the product. This product is not to be used by patients with kidney disease.

WARNINGS: Sulfonamides are known to cause Stevens-Johnson syndrome in hypersensitive individuals. Stevens-Johnson syndrome also has been reported following the use of sodium sulfacetamide topically. Cases of drug-induced systemic lupus erythematosus from topical sulfacetamide also have been reported. In one of these cases, there was a fatal outcome. **KEEP OUT OF THE REACH OF CHILDREN.**

PRECAUTIONS: FOR EXTERNAL USE ONLY. NOT FOR OPHTHALMIC USE.

General: Nonsusceptible organisms, including fungi, may proliferate with the use of this preparation.

Although rare, sensitivity to sodium sulfacetamide may occur. Therefore, caution and careful supervision should be observed when prescribing this drug for patients who may be prone to hypersensitivity to topical sulfonamides. If the use of this product produces signs of hypersensitivity or other untoward reactions, discontinue use of the preparation. Patients should be carefully observed for possible local irritation or sensitization during long-term therapy. Systemic toxic reactions such as agranulocytosis, acute hemolytic anemia, purpura hemorrhagica, drug fever, jaundice and contact dermatitis indicate hypersensitivity to sulfonamides. Particular caution should be employed if areas of denuded or abraded skin are involved. Systemic absorption of topical sulfonamides is greater following application to large, infected, abraded, denuded or severely burned areas. Under these circumstances, any of the adverse effects produced by the systemic administration of these agents could potentially occur, and appropriate observations and laboratory determinations should be performed.

Information for Patients: Patients should discontinue the use of this product if the condition becomes worse or if a rash develops in the area being treated or elsewhere. The use of this product also should be

discontinued promptly and the physician notified if any arthritis, fever or sores in the mouth develop. Avoid contact with eyes, lips and mucous membranes.

Drug Interactions: This product is incompatible with silver preparations.

Carcinogenesis, Mutagenesis and Impairment of Fertility:

Long-term animal studies for carcinogenic potential have not been performed on this product to date. Studies on reproduction and fertility also have not been performed. Chromosomal nondisjunction has been reported in the yeast, *Saccharomyces cerevisiae*, following application of sodium sulfacetamide. The significance of this finding to the topical use of sodium sulfacetamide in the human is unknown.

Pregnancy: *Category C.* Animal reproduction studies have not been conducted with this product. It is also not known whether this product can affect reproduction capacity or cause fetal harm when administered to a pregnant woman. This product should be used by a pregnant woman only if clearly needed or when potential benefits outweigh potential hazards to the fetus.

Nursing Mothers: It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when this product is administered to a nursing woman.

Pediatric Use: Safety and effectiveness in children under the age of 12 years have not been established.

ADVERSE REACTIONS: Reports of irritation and hypersensitivity to sodium sulfacetamide are uncommon. The following adverse reactions, reported after administration of sterile ophthalmic sodium sulfacetamide, are noteworthy: instances of Stevens-Johnson syndrome and instances of local hypersensitivity which progressed to a syndrome resembling systemic lupus erythematosus; in one case a fatal outcome was reported (see WARNINGS).

OVERDOSAGE: The oral LD₅₀ of sulfacetamide in mice is 16.5 g/kg. In the event of overdosage, emergency treatment should be started immediately.

Manifestations: Overdosage may cause nausea and vomiting. Large oral overdosage may cause hematuria, crystalluria and renal shutdown due to the precipitation of sulfa crystals in the renal tubules and the urinary tract. For treatment, contact your local

Poison Control Center or your doctor.

DOSAGE AND ADMINISTRATION:

WASH-OFF APPLICATION: Shake well before use. Cleanse affected skin thoroughly and pat dry before each application. Holding can upright, dispense product into palm of hand. Massage the product into the affected area and wait 10 minutes. Rinse thoroughly with water and pat dry. Treat the affected area 1 to 3 times daily, or as directed by a physician.

LEAVE-ON APPLICATION: Shake well before use. Cleanse affected skin thoroughly and pat dry before each application. Holding can upright, dispense product into palm of hand. Massage the product into the affected area. Wipe off any excess. Treat the affected area 1 to 3 times daily, or as directed by a physician.

STORAGE: Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F). Brief exposure to temperatures up to 40°C (104°F) may be tolerated provided the mean kinetic temperature does not exceed 25°C (77°F); however, such exposure should be minimized. Contents under pressure. Do not puncture or incinerate.

NOTICE: Protect from freezing and excessive heat. The product may tend to darken slightly on storage. Slight discoloration does not impair the efficacy or safety of the product. Keep dispensing container tightly closed.

Occasionally, a slight discoloration of fabric may occur when an excessive amount of the product is used and comes in contact with white fabrics. This discoloration, however, presents no problem, as it is readily removed by ordinary laundering without bleaches.

HOW SUPPLIED: This product is supplied in the following size(s):

3.5 oz. (100 g) can, **NDC** 0178-0696-01

To report a serious adverse event or obtain product information, call 1-800-298-1087.



Manufactured for:
MISSION PHARMACAL COMPANY
San Antonio, TX 78230 1355

OVA027R0616